Kinetic visual acuity is correlated with functional visual acuity at higher speeds

Ikko Iehisa, Kazuno Negishi , Masahiko Ayaki , Kazuo Tsubota

To cite: Iehisa I, Negishi K, Ayaki M, *et al*. Kinetic visual acuity is correlated with functional visual acuity at higher speeds. *BMJ Open Ophthalmology* 2019;4:e000383. doi:10.1136/bmjophth-2019-000383

II and KN contributed equally.

Received 13 August 2019
Revised 1 November 2019
Accepted 4 November 2019

ABSTRACT

Objective To measure the kinetic visual acuity (KVA) which is the ability to identify approaching objects and the functional visual acuity (FVA) which is continuous VA during 1 min under binocular and monocular condition (non-dominant eye shielding) for healthy subjects, and related ocular parameters to explore their correlation and implication in aspect of integrated visual function.

Methods The mean age of the 28 participants was 38.6±8.9 years (range, 23–57 years; 6 women). A KVA metre (AS-4Fα) and FVA metre (AS-28) were used to measure KVA and FVA, respectively. Multiple regression analysis was conducted to explore correlations among the measured visual function and related parameters, including age, binocularity, best-corrected visual acuity, refraction and tear break-up time.

Results The results of binocular KVA were better than monocular KVA at all speeds. A strong correlation was found between monocular and binocular KVA. The results of binocular FVA were better than monocular FVA ($p<0.001$) and there was a correlation between monocular and binocular FVA ($R=0.638$, $p<0.001$), as well as the maintenance rate for FVA ($R=0.228$, $p=0.003$). A linear mixed-effects model revealed that binocularity for KVA prediction was significant at all speeds and FVA was also significant at 60 km/h ($p<0.05$).

Conclusion The current results suggest that both binocularity and FVA may contribute to KVA.

Trial registration number UMIN00032385

INTRODUCTION

A better understanding of binocular visual function is required in modern society. People with monocular vision may suffer impaired visual function as the superiority of binocularity is well established in visual acuity (VA), reading speed, depth perception and movement detection.^{1–6} Binocular summation/binocularity is composed of probability summation and neural summation. When the light stimulates the photoreceptors in one eye, the corresponding photoreceptors in the contralateral eye are simultaneously stimulated. Thus, the probability of photoreceptor stimulation in the binocular conditions is higher with binocular than monocular viewing.³ Other studies reported that binocular viewing lowers the contrast threshold by up to 40%^{7,8} and the degree of summation is related to the complexity of the visual task.⁹

Key messages

What is already known about this subject?

- ▶ Previous investigations described binocular kinetic visual acuity (KVA) was superior to monocular KVA, however, there have been only a few investigations on other factors influencing differences in KVA.

What are the new findings?

- ▶ KVA was closely correlated with functional visual acuity (FVA) at higher speeds and this was stronger than the correlation between KVA and binocularity.

How might these results change the focus of research or clinical practice?

- ▶ Close correlation between KVA and FVA suggests that FVA could be a potential indicator for evaluating visual function in daily life to percept moving objects.

However, the functional role of binocularity has not been fully determined for daily activities. Common vision-threatening diseases that may cause monocular status include cataract, glaucoma and age-related macular degeneration, along with disorders of the central nervous system, such as cerebral infarction. A recent increase in the prevalence of the above-mentioned age-related eye diseases^{10–13} may be associated with increased traffic fatalities involving the elderly as binocular or monocular diseases can affect the integrated visual function essential for traffic safety. It is an emerging issue in our ageing society.

Kinetic VA (KVA) is the ability to identify approaching objects, whereas the ability to identify objects moving horizontally or vertically is called dynamic VA.¹⁴ There have been only a few investigations of KVA, and detailed studies have not been conducted on factors influencing differences in KVA ability. Rose³ reported that the threshold of movement detection for binocular viewing is quite small compared with monocular viewing, and the threshold for monocular viewing is higher than that for binocular viewing out to distances of 15–20 m, beyond which they are the same. This finding indicates that binocularity may contribute more at higher speeds to recognise distant objects.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Department of Ophthalmology, Keio University School of Medicine, Tokyo, Japan

Correspondence to

Dr Masahiko Ayaki; mayaki@olive.ocn.ne.jp

Table 1 Visual acuity and refraction of the participants

Visual acuity	Binocular	Monocular		P value (binocular vs dominant)	P value (binocular vs non-dominant)
		Dominant eye	Non-dominant eye		
UCVA (LogMAR)		0.49±0.56	0.50±0.57		
BCVA (LogMAR)	-0.20±0.06	-0.21±0.17	-0.16±0.06	0.493	0.021
SE (diopter)		-2.23±2.49	-3.06±3.72		

Data are presented as mean±SD.

BCVA, best corrected visual acuity; SE, spherical equivalent; UCVA, uncorrected visual acuity.

Recent investigations suggest that dry eye (DE) affects numerous visual parameters including higher-order aberration, accommodation and functional VA (FVA).^{15 16} DE is a very common eye disease and the estimated prevalence is 10%–20% in the adult population.¹⁷ DE is diagnosed by a specific corneal examination, lacrimal examination and medical interview.¹⁸ FVA is worse in DE and many other age-related eye diseases.¹⁹ The previous investigation described that among various clinical factors, FVA significantly predicted the correct answer rate in driving aptitude tests, and it was a promising method for screening driving aptitude, including both visual and cognitive functions in a short time.²⁰ However, the association between KVA and FVA has not been determined even though both are involved in the performance of watching objects continuously and are critical in driving safety.

KVA and FVA were examined under binocular and monocular conditions and analysed the correlation between these visual functions and ocular parameters. The aim of this study was to explore these correlations and find an appropriate method for determining the best visual performance with integrated visual function.

METHODS

Participants

Participants aged 20 or older were recruited via an advertisement in our research centre. The exclusion criteria were best-corrected visual acuity (BCVA) <20/30 in either eye, systemic disease or eye disease affecting visual function, history of surgery or refractive surgery, near

stereovision over 80s and a Mini-Mental State Examination (MMSE) score <24.

Ethics statement

The Institutional Review Board and Ethics Committee of the Keio University School of Medicine approved this study (approval number: 20170295) and the methods were carried out in accordance with the Declaration of Helsinki. Written informed consent was obtained from all participants. The study protocol was registered with the UMIN Clinical Trials Registry (UMIN00032385).

Ophthalmological and cognitive examination

Bilateral monocular and binocular distance VA was measured with the best corrective lenses (BCVA) and without corrective lenses (uncorrected VA). All participants underwent measurement of binocular-corrected and monocular-corrected distant FVA using the AS-28 FVA measurement system (Kowa, Aichi, Japan). The FVA was defined as the average of all VA values measured during 60s with the best distance correction. Measurements started with the Landolt ring size of the subject's BCVA. Each Landolt ring was displayed in the device for 2s. When the subject's response was correct, the next Landolt ring size became smaller. If the response was incorrect or none, the next Landolt ring size became larger. The maintenance rate of FVA was calculated as follows: (lowest logarithm of the minimum angle of resolution (logMAR) VA score – FVA at 60s) / (lowest logMAR VA score – baseline VA).

The KVA was measured using a KVA metre (AS-4; Kowa, Tokyo, Japan.) with the best distant correction as follows. The participants were instructed to look through the measuring window. They would detect the Landolt ring of 20/60 in a bright white circular shape. The Landolt ring moved towards the subjects from a 50m distance at a velocity of 20, 40 and 60km/h. The participants were asked to immediately respond by pressing a joystick when they identified the orientation of the Landolt ring. The results were expressed as decimal VA according to the distance of the participant's response. After two practices to reduce mistakes due to unfamiliarity with the test, the participants carried out five trials at three velocities under binocular and monocular conditions, with a 1 min rest between each setting. If the participant's answer was incorrect in all five trials, it was excluded from the analysis.

Table 2 Comparison of binocular and monocular kinetic visual acuity

Speed of the target (km/h)	Binocular	Monocular	P value*	Correlation between binocular and monocular†
20	0.22±0.14	0.29±0.13	<0.001	R=0.715, p<0.001
40	0.30±0.16	0.44±0.21	<0.001	R=0.755, p<0.001
60	0.38±0.19	0.52±0.23	<0.001	R=0.694, p<0.001

Data are mean±SD.

*Wilcoxon signed-rank test,

†Pearson's correlation coefficient.

Table 3 Simple linear regression analysis of kinetic visual acuity

Independent value	Correlation coefficient	P value
Speed of the target	0.391	<0.001
Binocular, monocular	-0.278	<0.001
UCVA in dominant eye	0.287	<0.001
UCVA in non-dominant eye	0.238	0.002
BCVA in dominant eye	0.204	0.008
BCVA in non-dominant eye	0.274	<0.001
SE in dominant eye	-0.241	0.002
SE in non-dominant eye	-0.206	0.007
FVA	0.473	<0.001
Maintenance rate of FVA	0.373	<0.001
Tear break up time (s)	-0.071	0.358

BCVA, best-corrected visual acuity; FVA, functional visual acuity; SE, spherical equivalent; UCVA, uncorrected visual acuity.

The stereo fly test (Stereo Optical Company Inc., Chicago, Illinois, USA) was performed at 40 cm to evaluate near stereopsis. The dominant eye was determined by the Rosenbach test. The Rosenbach method²¹ was used to determine which was the dominant eye. Subjects

concealed small objects far away with their thumbs under binocular vision. When one of the eyes closed, the object hidden by the thumb became visible, which proved that the closed eye was the dominant eye.

The standard tear BUT was measured after the instillation of fluorescein sodium in the conjunctival sac with a test filter paper.²² The interval between the last complete blink and the appearance of the first corneal black spot in the stained tear film was measured three times, and the mean value was calculated. The cut-off value was 5/6s. Cognitive status was evaluated with the MMSE-Japanese (MMSE-J, Nihon Bunka Kagakusha Co. Ltd, Tokyo, Japan).

Statistical analysis

For statistical analyses, decimal VA values were converted to the logarithm of the minimum angle of resolution. The measured value in the dominant eye was used for comparison with the binocular value. Where appropriate, data are expressed as the mean±SD. The normality of distribution of all variables was confirmed using the Shapiro-Wilk test with a significance level of 0.05. Paired *t* tests and Wilcoxon signed-rank tests were used for univariate analyses of continuous demographic and clinical features to compare monocular and binocular KVA and FVA. To further evaluate the association between

Figure 1 Scatterplot of KVA and FVA at each speed. There are significant correlations between KVA and FVA at 40 and 60 km/h: (A) Binocular, 20 km/h; (B) binocular, 40 km/h; (C) binocular, 60 km/h; (D) monocular, 20 km/h; (E) monocular, 40 km/h; (F) monocular, 60 km/h. FVA, functional visual acuity; KVA, kinetic visual acuity.

Table 4 Multiple regression analysis of kinetic visual acuity

Independent value	Non-standardised regression coefficient	SE	Normalised regression coefficient	P value	Collinearity test
Speed of the target	0.100	0.016	0.383	<0.001	1.000
BCVA in dominant eye	0.299	0.074	0.242	<0.001	1.008
FVA in non-dominant eye	1.340	0.163	0.493	<0.001	1.008

BCVA, best-corrected visual acuity; FVA, functional visual acuity.

KVA and other parameters, a linear mixed-effects model analysis was conducted with monocular and binocular KVA as dependent variables and demographic (age and sex) and ophthalmic parameters (monocular and binocular BCVA and FVA, FVA maintenance rate, refraction, anisometropia, BUT, Schirmer test value and MMSE score) as independent variables to obtain standardised correlation coefficients. All statistical tests were two-sided and the significance level was set to $\alpha=0.05$. All analyses were performed using SPSS V.24 software (IBM, Armonk, New York, USA).

Patient and public involvement

Study participants were involved in this study as the examinee. They were recruited via an advertisement in our research centre and fully informed of the time, burden and safety of examinations. The Institutional Review Board and Ethics Committee of the Keio University School of Medicine approved this study and were to regularly check the compliance.

RESULTS

This study included 28 volunteers (6 women and 22 men) with the mean age of 38.6 ± 8.9 years (range, 22–56 years). All participants satisfied the inclusion and exclusion criteria. The results of VA and refraction are shown in table 1. Most participants were mild myopes. The mean tear break-up time (BUT) was 9.64 ± 2.84 s and three participants had a short BUT and were possible patients with DE according to the latest definition.¹⁸

In all participants, near stereovision was over 80 s and the MMSE score was better than 24 points with a mean MMSE score of 29.9 ± 0.26 .

Comparisons of KVA between binocular and monocular conditions are shown in table 2. Binocular KVA was better than monocular at all speeds and a strong correlation was observed between binocular and monocular KVA. KVA at 60 km/h was significantly worse than KVA at 20 km/h under binocular and monocular conditions ($p<0.01$).

The results of a simple regression analysis between KVA and independent values (VA, refraction, FVA, object speed and binocular/monocular condition) are shown in table 3. The correlation between KVA and FVA ($R=0.473$, $p<0.001$) was stronger than between KVA and BCVA of the non-dominant eye ($R=0.274$, $p<0.001$) or binocularity ($R=0.278$, $p<0.001$). There were significant correlations

between KVA and FVA at 40 and 60 km/h under both binocular and monocular conditions (figure 1). BUT was not correlated with KVA at any speed. The results of multiple regression analysis between KVA and visual parameters are shown in table 4. The correlation between KVA and FVA in the non-dominant eye ($R=0.493$, $p<0.001$) was stronger than for BCVA in the dominant eye ($R=0.242$, $p<0.001$). The results of the analysis of factors affecting KVA using a linear mixed-effects model are shown in table 5. The standardised regression coefficient for binocularity was significant at all speeds and that of FVA was significant at 60 km/h (figure 2).

Binocular FVA (-0.13 ± 0.05) was significantly better than monocular FVA (-0.07 ± 0.09 ; Wilcoxon signed-rank tests, $p<0.001$), and a strong correlation existed between binocular and monocular FVA ($R=0.659$, $p<0.001$; figure 3).

DISCUSSION

The effects of binocularity and ocular parameters on KVA were evaluated and binocular FVA was correlated with KVA at higher speeds. It is speculated that both KVA and FVA are important for major integrated visual function working with continuous fixation. FVA may deteriorate in cataract, posterior capsule opacification, DE and age-related macular degeneration.^{23–28} These are treatable eye diseases and adequate ophthalmological treatments are recommended for drivers suffering these diseases to achieve the best possible visual performance. It is notable that FVA was more strongly correlated with KVA at a higher speed than age. Given that visual function is critical in driving, as well as physical mobility and recognition, vision care specialists and drivers should be aware that FVA provides the ability to keep seeing and focusing on objects.¹⁷

Although DE is a very common ocular surface disease and major cause of decreased FVA, which should be maintained in high-speed driving, the patients may be asymptomatic and diagnosed with symptoms as well as loss of homeostasis of the tear film.²⁹ Visual disturbance in patients with DE may be due to a blurry optical image with increased aberration and scattering induced by an unstable tear film.^{30 31} Patients with DE may recognise their symptoms only as eye fatigue or presbyopia, so DE may be often overlooked.³² KVA was correlated more strongly with FVA than BCVA and previous investigation

Figure 2 Standardised regression coefficient values for kinetic visual acuity, using a linear mixed-effects model. Binocularity was significant at all target speeds and FVA was significant at 60 km/h (* $p < 0.05$, linear mixed-effects model). FVA, functional visual acuity.

described a significant correlation between FVA and the useful field of view, which is related to the driver's cognitive function; therefore, FVA is an important indicator for driving safety, whereas KVA is important in recognising pedestrians and other traffic circumstances.²⁰ Based on the present results, the FVA examination could be

Figure 3 Scatterplot of binocular and monocular FVA. Binocular FVA was better than monocular FVA (Wilcoxon signed-rank tests; $p < 0.001$), and a strong correlation was found between the two parameters ($R = 0.659$, $p < 0.001$). FVA, functional visual acuity.

recommended in the driving aptitude test, as currently, only BCVA is required in Japan. FVA can be screened by a user-friendly automated instrument on a self-check basis, as used in the present study.

Binocular KVA was better than monocular KVA as expected. Many studies investigating the effect of binocularity report that visual function, including VA, speed cognition and contrast sensitivity, improve with binocular summation.^{1–3 7–9 33–35} For example, the visual performance of Formula one drivers, for example, sense of

Table 5 Linear mixed-effects model analysis results for factors affecting kinetic visual acuity

Independent variable	Regression coefficient	SE	Standardised regression coefficient	F value	P value
Speed: 20 km/h					
Binocularity*	−0.053	0.024	2.168	4.700	0.031
Sex†	0.110	0.070	1.574	2.476	0.129
FVA	0.184	0.412	0.446	0.199	0.656
Maintenance rate of FVA	−0.515	0.679	0.759	0.576	0.449
Age (y)	−0.002	0.003	0.635	0.403	0.532
Speed: 40 km/h					
Binocularity*	−0.077	0.026	2.968	8.810	0.003
Sex†	−0.051	0.084	0.608	0.370	0.549
FVA	0.507	0.451	1.126	1.268	0.262
Maintenance rate of FVA	−0.713	0.747	0.955	0.912	0.341
Age (y)	0.000	0.004	0.097	0.009	0.924
Speed: 60 km/h					
Binocularity*	−0.102	0.031	3.301	10.898	0.001
Sex†	0.028	0.095	0.297	0.088	0.770
FVA	1.267	0.547	2.317	5.370	0.022
Maintenance rate of FVA	1.278	0.880	1.452	2.109	0.148
Age (y)	0.002	0.004	0.539	0.291	0.595

*Binocular=1, monocular=0.

†Male=1, female=0.

FVA, functional visual acuity.

speed and distance, improves with a binocular summation.³⁵ The present results could provide additional evidence by showing the additive effects of binocularity in KVA and FVA. The present results also indicate that KVA is more strongly correlated with BCVA in the non-dominant eye than with BCVA in the dominant eye, suggesting that the non-dominant eye may significantly contribute to binocularity with appropriate refractive correction.

The limitations of the present study first include the small number of participants. Second, the significance of FVA should be confirmed with a larger number of patients with severe DE, before and after the intervention. Third, potential contribution with adaptation was not excluded in the verification of the superiority of binocular vision in the present study. Thus, it should be claimed as another limitation of our study as the results of the binocular test may be overestimated. Finally, the present results should be further evaluated in older people with impaired MMSE and visual function to predict the risk of motor accidents potentially caused by insufficient visual performance.

CONCLUSIONS

BCVA, KVA and FVA were strongly correlated with each other and FVA was contributory to KVA at high speed. The present results may be applicable for the evaluation of driving aptitude in diseased and older individuals.

Contributors II: collected and analyzed the data. II and MA: wrote the manuscript. II and KN: designed the study. II, MA, KN and KT: interpreted the data for the work, reviewed and approved the final version of the manuscript.

Funding Hitachi Automotive Systems, Ltd.

Disclaimer II and KN work have been funded by Hitachi Automotive Systems, Ltd. KT has the patent (US patent No.7470026) of kinetic visual acuity meter (AS-4Fo).

Competing interests None declared.

Patient consent for publication Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon request

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Kazuno Negishi <http://orcid.org/0000-0002-2892-9810>

Masahiko Ayaki <http://orcid.org/0000-0002-9481-0164>

REFERENCES

- Horowitz MW. An analysis of the superiority of binocular over monocular visual acuity. *J Exp Psychol* 1949;39:581–96.
- Jones RK, Lee DN. Why two eyes are better than one: the two views of binocular vision. *J Exp Psychol* 1981;7:30–40.
- Rose D. Monocular versus binocular contrast thresholds for movement and pattern. *Perception* 1978;7:195–200.
- McCann BC, Hayhoe MM, Geisler WS. Contributions of monocular and binocular cues to distance discrimination in natural scenes. *J Vis* 2018;18:12.
- Yu D, Cheung S-H, Legge GE, et al. Reading speed in the peripheral visual field of older adults: does it benefit from perceptual learning? *Vision Res* 2010;50:860–9.
- Johansson J, Pansell T, Ygge J, et al. Monocular and binocular reading performance in subjects with normal binocular vision. *Clin Exp Optom* 2014;97:n/a–8.
- Campbell FW, Green DG. Monocular versus binocular visual acuity. *Nature* 1965;208:191–2.
- Wehrhahn C, Westheimer G, Abulencia A. Binocular summation in temporal-order detection. *J Opt Soc Am A* 1990;7:731–2.
- Frisén L, Lindblom B. Binocular summation in humans: evidence for a hierarchic model. *J Physiol* 1988;402:773–82.
- Agramunt S, Meuleners LB, Fraser ML, et al. Bilateral cataract, crash risk, driving performance, and self-regulation practices among older drivers. *J Cataract Refract Surg* 2016;42:788–94.
- Kwon M, Huisingh C, Rhodes LA, et al. Association between glaucoma and At-fault motor vehicle collision involvement among older drivers. *Ophthalmology* 2016;123:109–16.
- Wood JM, Black AA, Mallon K, et al. Effects of age-related macular degeneration on driving performance. *Invest Ophthalmol Vis Sci* 2018;59:273–9.
- Huisingh C, McGwin G, Wood J, et al. The driving visual field and a history of motor vehicle collision involvement in older drivers: a population-based examination. *Invest Ophthalmol Vis Sci* 2015;56:132–8.
- Hoshina K, Tagami Y, Mimura O, et al. A study of static, kinetic, and dynamic visual acuity in 102 Japanese professional baseball players. *OPHTH* 2013;7:627–32.
- Goto E, Yagi Y, Matsumoto Y, et al. Impaired functional visual acuity of dry eye patients. *Am J Ophthalmol* 2002;133:181–6.
- Goto E, Yagi Y, Kaido M, et al. Improved functional visual acuity after punctal occlusion in dry eye patients. *Am J Ophthalmol* 2003;135:704–5.
- Uchino M, Nishiwaki Y, Michikawa T, et al. Prevalence and risk factors of dry eye disease in Japan: Koumi study. *Ophthalmology* 2011;118:2361–7.
- Tsubota K, Yokoi N, Shimazaki J, et al. New perspectives on dry eye definition and diagnosis: a consensus report by the Asia dry eye Society. *Ocul Surf* 2017;15:65–76.
- Kaido M, Kawashima M, Ishida R, et al. Severe symptoms of short tear break-up time dry eye are associated with accommodative microfluctuations. *Clin Ophthalmol* 2017;11:861–9.
- Negishi K, Masui S, Mimura M, et al. Relationship between functional visual acuity and useful field of view in elderly drivers. *PLoS One* 2016;11:e0147516.
- Rosenbach O. Über monokulare Vorherrschaft beim binocularen Sehen. *Med Wochenschr* 1903;50:1290–2.
- Yokoi N, Georgiev GA, Kato H, et al. Classification of Fluorescein Breakup Patterns: A Novel Method of Differential Diagnosis for Dry Eye. *Am J Ophthalmol* 2017;180:72–85.
- Yamaguchi T, Negishi K, Tsubota K. Functional visual acuity measurement in cataract and intraocular lens implantation. *Curr Opin Ophthalmol* 2011;22:31–6.
- Yamaguchi T, Negishi K, Dogru M, et al. Improvement of functional visual acuity after cataract surgery in patients with good pre- and postoperative Spectacle-Corrected visual acuity. *J Refract Surg* 2009;25:410–5.
- Watanabe K, Negishi K, Kawai M, et al. Effect of experimentally induced astigmatism on functional, conventional, and low-contrast visual acuity. *J Refract Surg* 2013;29:19–25.
- Katada Y, Negishi K, Watanabe K, et al. Functional visual acuity of early presbyopia. *PLoS One* 2016;11:e0151094.
- Yotsukura E, Torii H, Saiki M, et al. Effect of neodymium:YAG laser capsulotomy on visual function in patients with posterior capsule opacification and good visual acuity. *J Cataract Refract Surg* 2016;42:399–404.
- Tomita Y, Nagai N, Suzuki M, et al. Functional visual acuity in age-related macular degeneration. *Optom Vis Sci* 2016;93:70–6.
- Craig JP, Nichols KK, Akpek EK, et al. TFOS DEWS II definition and classification report. *Ocul Surf* 2017;15:276–83.
- Kaido M, Toda I, Oobayashi T, et al. Reducing short-wavelength blue light in dry eye patients with unstable tear film improves performance on tests of visual acuity. *PLoS One* 2016;11:e0152936.
- Koh S, Tung CI, Inoue Y, et al. Effects of tear film dynamics on quality of vision. *Br J Ophthalmol* 2018;102:1615–20.
- Toda I, Fujishima H, Tsubota K. Ocular fatigue is the major symptom of dry eye. *Acta Ophthalmol* 1993;71:347–52.
- Wardle SG, Alais D. Evidence for speed sensitivity to motion in depth from binocular cues. *J Vis* 2013;13:17.
- Lema SA, Blake R. Binocular summation in normal and stereoblind humans. *Vision Res* 1977;17:691–5.
- Westlake W. Is a one eyed racing driver safe to compete? formula one (eye) or two? *Br J Ophthalmol* 2001;85:619–24.