

Economic burden and loss of quality of life from dry eye disease in Canada

 Clara Chan ,¹ Setareh Ziai,² Varun Myageri,³ James G Burns,⁴ C Lisa Prokopich⁵

To cite: Chan C, Ziai S, Myageri V, *et al.* Economic burden and loss of quality of life from dry eye disease in Canada. *BMJ Open Ophthalmology* 2021;**6**:e000709. doi:10.1136/bmjophth-2021-000709

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjophth-2021-000709>).

Received 6 January 2021
Accepted 3 September 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Ophthalmology and Vision Sciences, University of Toronto, Toronto, Ontario, Canada

²University of Ottawa Eye Institute, Ottawa, Ontario, Canada

³IQVIA Solutions Canada Inc, Kirkland, Quebec, Canada

⁴Novartis Pharmaceuticals Canada Inc, Dorval, Quebec, Canada

⁵Optometry & Vision Science, University of Waterloo, Waterloo, Ontario, Canada

Correspondence to

Dr Clara Chan; clara.chan@gmail.com

ABSTRACT

Objective To describe the direct and indirect cost estimates of dry eye disease (DED), stratified by disease severity, and the impact of DED on quality of life (QoL) in Canadian patients.

Methods and analysis A prospective, multicentre, observational, cross-sectional study was conducted at six sites across Canada. Eligible patients completed a 20 min survey on demography, general health, disease severity, QoL and direct (resource utilisation and out-of-pocket expenses for the past 3–24 months) and indirect costs (absenteeism and presenteeism based on Work Productivity and Activity Impairment questionnaire responses). Subgroup analyses were performed according to DED severity and presence of Sjögren's syndrome.

Results Responses from 146 of 151 participants were included in the analysis. DED was rated as moderate or severe by 19.2% and 69.2% of patients, respectively. Total mean annual costs of DED were \$C24 331 (Canadian dollars) per patient and increased with patient-reported disease severity. Mean (standard deviation [SD]) indirect costs for mild, moderate and severe disease were \$C5961 (\$C6275), \$C16 525 (\$C11 607), and \$C25 485 (\$C22,879), respectively. Mean (SD) direct costs were \$C958 (\$C1216), \$C1303 (\$C1574) and \$C2766 (\$C7161), respectively. QoL scores were lowest in patients with Sjögren's syndrome (8.2% of cohort) and those with severe DED.

Conclusion This study provides important insights into the negative impact of DED in a Canadian setting. Severe DED was associated with higher direct and indirect costs and lower QoL compared with those with mild or moderate disease. Increased costs and poorer QoL were also evident for patients with DED plus Sjögren's syndrome versus DED alone.

INTRODUCTION

Dry eye disease (DED) is a complex, multifactorial disorder of the periocular tear film that may result in damage to the ocular surface and is associated with symptoms of ocular discomfort.¹ Globally, the estimated prevalence of DED ranges between 5% and 50%.² In Canada, the prevalence of DED has been estimated at 21%.³

Diagnosis of DED should consider symptomatology, visual disturbance, tear film stability and composition, tear volume, damage and/or inflammation of the ocular surface and eyelid aspects (eg, blepharitis, lid wiper epitheliopathy).^{4 5} In particular, the

Key messages

What is already known about this subject?

► Dry eye disease (DED) is associated with high direct and indirect socioeconomic costs as well as a considerable reduction in the patient's quality of life (QoL). However, there is a paucity of these data for DED in the Canadian healthcare system.

What are the new findings?

► This multisite study provides further data and confirms the particular socioeconomic and QoL impacts on Canadian patients with DED. We found that indirect costs to Canadian patients represent the largest proportion of overall costs.

How might these results change the focus of research or clinical practice?

► These data support the importance of effective management of DED to offset the substantial indirect costs and QoL burden on patients. Further research should focus on more detailed economic and QoL implications as well as the impact of pharmacological therapies and other interventions such as air quality adjustments or regular work breaks.

major cause of DED needs to be recognised before treatment plans can be developed.

The Tear Film and Ocular Surface Society (TFOS) Dry Eye Workshop (DEWS) II proposed a stepwise treatment approach with follow-up to monitor signs and symptoms of DED, in order to ensure that other comorbid ocular surface diseases are also treated appropriately.^{1 6} Initially, DED is often treated with ocular lubricants. If this approach proves inadequate, prescribed topical drugs (eg, antibiotics, corticosteroids, secretagogues and immunomodulatory or lymphocyte function-associated antigen-1 antagonist drugs) or oral antibiotics can be considered. Additional options include overnight chamber devices and tear conservation measures.⁶ Where treatment remains ineffective, the next step is to use serum eye drops, therapeutic contact lenses or oral secretagogues. If the above measures fail to control the disease, the TFOS DEWS II report identifies further options as lacrimal duct occlusion or longer term topical corticosteroids (monitored closely to

minimise potential adverse side effects such as cataract formation and glaucoma).⁶

DED can incur very high socioeconomic costs, both direct (eg, medical fees, prescribed and over-the-counter drugs) and indirect (eg, unemployment, work absenteeism, presenteeism (at work but impaired productivity)).⁷ Indirect costs comprise most of the overall economic burden. For example, in the USA, the mean average indirect cost to society was US\$55.4 billion compared with US\$3.8 billion for direct costs.⁸ Furthermore, DED can place a substantial burden on the quality of life (QoL) of patients, impacting their physical, social and psychological well-being and affecting their workplace productivity.^{7,9} For example, blurred and/or fluctuating vision may restrict activities of daily living such as reading, driving, watching television and using smartphones.⁷

The physical impact of DED pain can be likened to a type of chronic pain syndrome, negatively affecting patients' psychological and physical well-being and QoL.⁷ Several studies, as reviewed by McDonald *et al*,⁹ have demonstrated the impact that DED has on QoL. However, there are few publications describing the impact of DED on socioeconomic costs and QoL in Canada.^{2,3,10} The present study aimed to contribute to the literature on DED by capturing the direct and indirect cost estimates of DED, stratified by the severity of DED and to understand the impact on QoL for Canadian patients with DED.

MATERIALS AND METHODS

Study design

This was a prospective, multicentre, observational, cross-sectional study conducted at six optometry and ophthalmology sites across Canada. Eligible patients had a diagnosis of mild to severe DED by an accredited healthcare provider (HCP). The investigators prioritised patients with moderate to severe DED, and target recruitment for each site was set as 10%, 45% and 45% of patients with mild, moderate or severe DED, respectively, based on HCP classification.

Patients

Investigators recruited patients for the study according to their diagnosis of DED. Eligible patients were aged 18 to 64 years and were required to meet all of the following criteria: a current diagnosis of mild-to-severe DED made by an accredited healthcare professional; a routine visit to a recruiting healthcare professional; symptoms of dry eye for at least 1 year before the date of recruitment; literacy in English, ability to read and complete surveys in English and ability to read, understand and sign the informed consent form on a voluntary basis. Patients were excluded if they were already participating in another clinical trial.

Objectives

The two primary objectives of the study were to describe the direct out-of-pocket costs and indirect costs (absenteeism

and presenteeism) attributable to DED and to determine the QoL impact attributable to DED. The four secondary objectives comprised the above analyses conducted in groups stratified by patient-reported DED severity (mild, moderate or severe) and by Sjögren's syndrome status (diagnosed vs no known Sjögren's syndrome).

Data collection and assessments

All patients who provided consent were asked to complete a survey lasting approximately 20 min (on their own or with help) during their routine optometrist/ophthalmologist visit. Study data were collected from the survey, which was composed of six sections: demographic data, general health data, DED severity data, QoL data, indirect costs and direct costs (online supplemental table S1). Demographic data (age, sex, ethnicity, education, employment status) and clinical characteristics data (smoking status, contact lens use, screen exposure, DED treatment, DED risk factors, DED duration, ocular and non-ocular comorbidities) over the past 12 months were collected. Patient-reported DED severity and presence of Sjögren's syndrome were captured. Patients were asked whether they had been formally diagnosed with Sjögren's syndrome. Possible answers were 'yes', 'no', and 'I don't know'; surveys with this question left unanswered were listed as 'unknown'. Formal testing (eg, serology or minor salivary gland biopsy) was not performed to rule out Sjögren's syndrome in patients who did not receive a positive diagnosis.

Current DED severity was assessed according to patient self-rating of their discomfort with eye dryness using the Eye Dryness Score Visual Analog Scale (EDS VAS; score range 0–100 (no discomfort to maximum discomfort)). Scores of <40 were considered mild, 40 to <60 moderate and ≥60 severe DED. The EDS VAS has not been validated for the evaluation of DED severity; however, inclusion of a VAS as a key component of a DED questionnaire has been shown as having consistently good repeatability.¹¹

All costs are reported in 2018 Canadian Dollars. Annual direct costs were calculated using patient-reported resource utilisation and out-of-pocket expenses for the past 3–24 months. Five categories of treatments were used to assess out-of-pocket costs to patients with DED: ocular lubricants, cyclosporine, punctal plugs, optometrist or ophthalmologist visits and nutritional supplements. A patient's direct annual cost was the sum of their annualised cost types. Population and subgroup direct costs were calculated using the average of each patient's direct annual cost values.

Annual indirect costs were calculated with the human capital approach based on each patient's annual salary and their Work Productivity and Activity Impairment (WPAI) questionnaire¹² scores (online supplemental table S2). Average salary was estimated according to age group, sex, employment status and highest level of education from Statistics Canada,¹³ adjusted to 2018 levels using the Bank of Canada consumer price index inflation calculator.¹⁴ The WPAI questionnaire comprises

questions on how many hours at work were missed over the past 7 days due to DED (absenteeism) and the percentage of DED-related impairment in productivity while working over the past 7 days (presenteeism). The WPAI questionnaire has been validated to quantify work impairment for several diseases and to compare impairment between study treatment groups or individuals with different disease severity levels.^{15–21} It has also been used to evaluate impairment related to DED.^{22–24}

Current QoL was evaluated according to the National Eye Institute Visual Function Questionnaire (NEI VFQ 25).²⁵ The 25 questions of the VFQ were recoded into 12 subscale variables: general health, general vision, ocular pain, near activities, distance activities, social functioning, mental health, role difficulties, dependency, driving, colour vision and peripheral vision. For each subscale, numeric responses were converted according to scoring rules to values on a 0–100 scale, and the averages of the responses were calculated for each. Lower scores indicate poorer QoL. Overall QoL scores were calculated by averaging vision-targeted subscale scores, excluding general health.

Sample size and statistical analysis

The study aimed to recruit approximately 150 patients from six sites over 6 months; assuming a SD-to-mean ratio of 0.14, this would give a relative precision in the primary analysis of up to 7%, with the smallest cohort estimated to contain 15 surveys. This SD-to-mean ratio was derived from a 2013 US survey of patients with DED,²⁶ which determined mean and SD for Short Form-6 dimension (SF-6D) Health Utility Index results to be 0.7 and 0.1, respectively, representing an SD-to-mean ratio of 0.14 and a precision of 2.29%. According to Yu *et al*, the SD-to-mean cost ratio from the payer's perspective was closer to 0.1.⁸ For our sample size calculations, we assumed the ratio between SD and sample mean to be 0.1, resulting in a precision estimate of approximately 1.6% ($1.96 \times 0.1 \sqrt{150}$, where 1.96 represents the value of the 97.5th percentile of the normal distribution to calculate a two-sided 95% CI).

Data analysis was purely descriptive and this study did not explore any associations. Summary statistics for categorical variables included frequency and percentage of each category or modality. Summary statistics for continuous variables are reported as mean and SD.

RESULTS

Patients

In total, 151 patients from six Canadian optometrist and/or ophthalmologist sites consented to participate and the study lasted for 7 months (from August 2018 to March 2019). Data from 146 patients were analysed. Data from the remaining five patients were excluded from the analysis, mainly because they had not experienced DED symptoms for at least 1 year prior to recruitment. Demographic characteristics are shown in table 1. The mean (SD) age was 49.8 (11.4) years. Most patients were women

Table 1 Patient demographics

All values are n (%) unless stated otherwise	Patients (N=146)
Mean (SD) age, years	49.8 (11.4)
Female	131 (89.7)
Age group, years	
18–24	1 (0.7)
25–34	19 (13.0)
35–44	29 (19.9)
45–54	35 (24.0)
55–64	62 (42.5)
Ethnicity*	
Caucasian	104 (71.2)
East Asian	26 (17.8)
South Asian	8 (5.5)
Latino/Hispanic	3 (2.1)
African	3 (2.1)
Middle Eastern	2 (1.4)
Caribbean	2 (1.4)
Other†	4 (2.7)
Highest level of education	
University certificate, diploma or degree	
Above bachelor level	22 (15.1)
At bachelor level	45 (30.8)
Below bachelor level	4 (2.7)
College, CEGEP or other non-university certificate or diploma	34 (23.3)
Trades	
Certificate of apprenticeship or qualification	2 (1.4)
Certificate or diploma (excluding certificate of apprenticeship)	9 (6.2)
High school diploma or equivalency certificate	26 (17.8)
No certificate, diploma, or degree	4 (2.7)
Employment status	
Employed	95 (65.1)
Disability leave	
Due to DED	3 (2.1)
Other reason	9 (6.2)
Unemployed	
Job seeking	9 (6.2)
Not job seeking	9 (6.2)
Retired/pre-pension plan	15 (10.3)
None of the above ('I do something else')	6 (4.1)
Estimated income,‡ CAD	
Mean (SD)	\$C68 781 (\$C22 983)

Continued

Table 1 Continued

All values are n (%) unless stated otherwise	Patients (N=146)
Median (Q1, Q3)	\$C58 755 (\$C50 513, \$C87 415)
Estimated annual income,† CAD	
≤\$C40 000	4 (2.7)
>\$C40 000–≤\$C60 000	72 (49.3)
>\$C60 000–≤\$C80 000	24 (16.4)
>\$C80 000–≤\$C100 000	32 (21.9)
>\$C100 000	14 (9.6)

*As three patients were of mixed ethnicity the total does not add up to 100%.

†Includes one Filipino; the remaining three patients did not specify their ethnicity.

‡Average salary was estimated according to age group, sex, employment status, and highest level of education from Statistics Canada,¹³ adjusted to 2018 levels using the Bank of Canada consumer price index inflation calculator.¹⁴

CAD, Canadian dollars; CEGEP, College of General and Vocation Education; DED, dry eye disease.

(89.7%), of whom almost half (62/131 (47.3%)) were aged ≥55 years. The majority of patients were Caucasian (71.2%), followed by East Asians (17.8%), and there were only a few patients with mixed ethnicities (2.1%). Most patients (71.9%) held a college degree or a higher qualification, with approximately half of the group (45.9%) educated to university degree level (at or above bachelor level). The remaining patients (28.1%) did not hold a college or higher degree. Most patients were employed (65.1%) and only 2.1% were on disability leave as a result of DED. The estimated mean (SD) annual income was \$C68 781 (\$C22 983), with almost half (49.3%) of patients expected to earn >\$C40 000 but ≤\$C60 000.

Clinical characteristics are shown in table 2. Most patients (64.4%) had never smoked, 29.5% had smoked previously and only 6.2% were current smokers. A minority of patients used contact lenses (13.7%). Most patients (82.3%) spent more than 3 hours per day looking at screens, with 24.7% looking at screens for at least 7 hours per day. DED was rated on the EDS VAS as moderate or severe by 19.2% and 69.2% of patients, respectively. The analysis population only included patients who had been diagnosed with DED for at least 1 year: 90 patients (61.6%) had experienced DED for 1–5 years and the remainder had a disease duration of ≥6 years.

An absence of ocular comorbidities and/or surgeries/injections accounted for 52.7% and 67.8% of patients, respectively. The remainder reported a variety of ocular comorbidities, of which the most frequent were seasonal allergies with itchy eyes (21.9%) and Meibomian gland dysfunction (18.5%). The most frequent ocular surgery was refractive eye surgery, which had been conducted in 15.8% of patients. The most frequently used treatment reported by patients was eye drops (preserved,

Table 2 Clinical characteristics

All values are n (%) unless stated otherwise	Patients (N=146)
Use contact lenses	20 (13.7)
Screen exposure, hours	
0–<1	10 (6.8)
1–2	16 (11.0)
3–4	48 (32.9)
5–6	36 (24.7)
7–15	36 (24.7)
Current type of treatment	
Preserved	33 (22.6)
Non-preserved (eg, hydroxypropyl cellulose ophthalmic insert)	52 (35.6)
Prescription eye drops (eg, lifitegrast, cyclosporine)	73 (50.0)
Specialty drugs	11 (7.5)
Compounded ointments	12 (8.2)
Gel (eg, loteprednol etabonate)	21 (21.2)
Other	39 (26.7)
None	8 (5.5)
Sjögren's syndrome	
No	122 (83.6)
Yes	12 (8.2)
I do not know	11 (7.5)
Unknown ^b	1 (0.7)
Patient-reported severity of DED	
Mild	17 (11.6)
Moderate	28 (19.2)
Severe	101 (69.2)
DED duration, years	
1–5	90 (61.6)
6–10	28 (26.0)
10–35	18 (12.3)

*'Unknown' indicates that no answer was provided.

†Including hypertension, antidiabetic and analgesic medications.

‡For example, Sjögren's syndrome, rheumatoid arthritis, lupus.

§For example, mood, depression, anxiety.

¶Including laser-assisted in situ keratomileusis or photorefractive keratectomy.

DED, dry eye disease; GvHD, graft versus host disease; HRT, hormone replacement therapy; HSCT, haematopoietic stem cell transplantation.

22.6%; non-preserved, 35.6%). Concomitant medications included multivitamins, which were being taken by 40.4% of patients and antidepressants (used by 15.1% of patients). A total of 21.9% of patients were not taking any concomitant medications.

Twelve patients (8.2%) reported having Sjögren's syndrome and the remaining 134 (91.8%) had no known Sjögren's syndrome, including 122 (83.6%) who




Figure 1 Mean annual costs due to DED. CAD, Canadian dollars; DED, dry eye disease.

indicated that they had not received this diagnosis and 12 patients (8.2%) in whom the status was unknown. Among other diagnostic risk factors, thyroid disorders were reported by 15.1% of patients, whereas 32.2% did not report any diagnostic risk factors.

Annual costs of DED

The total annual costs of DED per patient were on average \$C24331 (figure 1); the majority was related to the indirect costs of DED presenteeism (79.3%). The mean (SD) patient-reported indirect cost attributable to DED was \$C21052 (\$C20812)/year (figure 1). Mean (SD) indirect costs of presenteeism and absenteeism were \$C19304 (\$C18916)/year and \$C2702 (\$C11028)/year, respectively. The mean (SD) total patient-reported direct cost attributable to DED was \$C2324 (\$6159)/year (figure 1).

When average annual costs were stratified by patient-reported DED severity, the mean (SD) indirect costs of absenteeism and presenteeism attributable to severe DED totaled \$C25485 (\$C22879)/year, representing an increase of 54% versus the costs of moderate DED (\$C16525 (\$C11607)/year) and 328% vs mild DED (\$C5961 (\$C6275)/year) (figure 2). Direct out-of-pocket costs per year were also highest for severe DED (mean (SD): \$C2766 (\$C7161)), representing an increase in 112% versus the costs of moderate DED (\$C1303 (\$C1574)) and 189% versus mild DED (\$C958 (\$C1216)).

The mean (SD) indirect costs per year of absenteeism and presenteeism attributable to Sjögren's syndrome




Figure 3 Mean annual costs according to Sjögren's syndrome status. CAD, Canadian dollars.

totalled \$C41094 (\$C15720), representing an increase in 132% versus the costs for patients with no known Sjögren's syndrome (\$C17694 (\$C17153)) (figure 3). Direct out-of-pocket costs were also higher for Sjögren's syndrome (mean (SD): \$C2689 (\$C2430)), representing an increase in 22% versus the costs for patients with no known Sjögren's syndrome (\$C2203 (\$C6660)).

Impact of DED on QoL

The total mean (SD) VFQ-25 QoL score was 77.13 (15.69) (figure 4). Ocular pain had the lowest score of all subscales (mean (SD) score 52.40 (21.99)), indicating that it had the greatest negative effect on QoL. DED also impacted mental health (66.14 (25.84)), role difficulties (at work and for other activities, 68.07 (26.47)) and driving (75.15 (19.38)).

On average, self-reported severe DED was associated with a lower total VFQ-25 QoL score (mean 72.42 (SD 16.19)) than moderate (86.60 (6.98)) or mild (89.55 (7.14)) DED. The mean (SD) total VFQ-25 QoL score was lower in patients with Sjögren's syndrome compared with those without the condition (59.81 (14.16) vs 79.21 (15.04)), respectively.




Figure 2 Mean annual costs according to patient-reported DED severity. CAD, Canadian dollars; DED, dry eye disease.




Figure 4 Average QoL subscale scores. Scores for the 12 VFQ 25 QoL subscales and the total QoL score, based on score ranges of 0–100, with lower scores indicating lower QoL. DED, dry eye disease; QoL, quality of life, VFQ, Visual Function Questionnaire.

DISCUSSION

This study provides an analysis of the economic burden and effects on QoL in Canadian patients with DED. Patients with self-reported severe DED reported lower QoL, higher medical fees and impaired work productivity compared with mild and moderate cases. Annual indirect costs increased from a mean (SD) of \$C5961 (\$C6275) in mild DED to \$C25 485 (\$C22 879) in severe DED. Direct costs followed a similar pattern, with a mean (SD) of \$C958 (\$C1216)/year in mild DED rising to \$C2766 (\$C7161)/year in severe DED. Patients with Sjögren's syndrome also incurred higher costs than those without Sjögren's syndrome. The mean total QoL score was 77, with ocular pain impacting patients' QoL the most (mean NEI VFQ-25 score 52); DED also negatively affected patients' mental health, role difficulties and driving. Unsurprisingly, QoL was worse in patients with severe DED versus those with mild/moderate disease and in those with versus those without Sjögren's syndrome.

Much of the previous literature have evaluated the costs and burden of DED from a payers' or healthcare systems perspective and have found that DED confers a substantial economic burden. In a systematic literature review, the majority of economic data was found to be from the USA (9/12 articles), and health-related QoL data were predominantly from Europe (11/20 articles) and the USA (8/20 articles).⁹ McDonald *et al* found that indirect costs comprised the largest proportion of overall DED costs, due to significant loss in work productivity.⁹ In the USA, for example, indirect costs were US\$11 302/year per patient (with the overall burden to society being US\$55 billion), and the numbers of days lost per year for patient-reported mild, moderate and severe DED were 91, 95 and 128, respectively.⁸ None of the 12 economic burden articles included Canadian data, although one article reported Canadian QoL data.

Our research provides additional insights into the burden of DED in a Canadian setting, taking the patients' direct costs into account and estimating indirect costs to society. We have shown that in Canada, indirect costs comprise the largest proportion of overall costs compared with direct costs. We found that DED impacts both indirect and direct costs. In addition, direct costs increased with disease severity. Our data are supported by an analysis of 2005 data from 2171 US patients, where indirect versus direct costs were estimated as US\$11 302/year versus US\$783/year per patient.⁸ Likewise, direct costs increased with patient-reported disease severity (US\$678/year for mild DED, US\$771 for moderate DED and US\$1267 for severe DED). In this study, both prescription and non-prescription medications were considered in the five categories of healthcare resource use (ocular lubricant treatment, cyclosporine, punctal plugs, physician visits and nutritional supplements).

The WPAI questionnaire is a validated questionnaire for the evaluation of work impairment.^{15–21} It was associated with higher construct validity and fewer omissions when administered by an interviewer rather than

self-administered.¹⁵ The WPAI was also found to provide higher mean estimates of productivity loss and related cost compared with other questionnaires.^{27 28} To evaluate ixekizumab therapy on work productivity among patients with chronic plaque psoriasis (PSO), Armstrong *et al* used the WPAI-PSO questionnaire.²⁹ Results were represented as observed mean percentages of absenteeism, presenteeism, work productivity loss (overall work impairment associated with absenteeism and presenteeism) and activity impairment, as derived from patient assessment of each factor on a scale of 0–10.

Direct costs determined in our study were considerably higher than those found in an analysis of 2003–2004 DED management data from six European countries (France, Germany, Italy, Spain, Sweden and the UK).³⁰ To treat 1000 patients with DED, estimated direct medical costs (patient examinations, diagnostic tests, prescription medications and surgical procedures) ranged from US\$270 000 (France) to US\$1.1 million (UK); the per-patient average amounts converted to 2018 Canadian dollars (estimated \$C1.13 conversion rate³¹ inflated with the Consumer Price Index¹⁴) would be approximately \$C360–\$1460. However, these figures reflect the data captured through management by an ophthalmologist only; self-treatment with over-the-counter agents and therapies prescribed by a general practitioner were not included.

Similar to our data, in which patients ranked ocular pain as the lowest favourable item in terms of QoL, impaired QoL in patients with active primary Sjögren's syndrome is mostly caused by ocular pain and dryness.³² In the USA, the mean ocular pain subscale score was significantly lower for patients with moderate-to-severe DED compared with patients with milder DED.³³ Additionally, assessment of the relative burden of DED in the USA showed that DED consistently caused bodily pain (effect size –0.08) and decreased role-physical (defined as limitations due to physical problems, –0.07) and vitality scores (–0.11) when compared with people without DED. However, the difference was only clinically significant for moderate or severe DED.³⁴ QoL was impaired to a greater extent in patients with Sjögren's syndrome versus those with non-Sjögren's syndrome DED.³³ Our data are in agreement with these observations.

Although outside the scope of this analysis, additional factors in the severity of DED should be taken into consideration. The overwhelming majority (89.7%) of patients were women, and nearly half (42.5%) of the patients in this study were aged 55 years or older. Both of these characteristics are significantly associated with not only increased prevalence but also greater severity of DED. According to a study by Tellefsen Nøland *et al*, women had significant increased osmolarity, shorter tear break-up time (TBUT), decreased meibum quality, reduced meibum expressibility and decreased corneal sensitivity than men.³⁵ Increased age was associated with significantly worse Ocular Surface Disease Index (OSDI) score, shorter TBUT, lower Schirmer I test score and

reduced meibum expressibility. Postmenopausal women were found to have higher scores related to ocular symptoms, vision-related functions and environmental triggers than perimenopausal women, and OSDI score increased with age.³⁶ Screen exposure has also been shown to have a significant effect on DED. One-half of subjects in this study reported at least 5 hours of daily screen time. Prolonged digital exposure time was a significant risk factor for DED in several studies,^{37–40} and significantly shorter TBUT times were observed in patients with ≥ 8 hours of daily screen time.⁴⁰ Smartphone usage was strongly associated with paediatric DED.⁴¹

While our findings provide important evidence on the economic and QoL burden associated with DED in a Canadian population, they must be interpreted in context. Canada has a distinct climate with the potential to worsen the symptoms of DED as a result of dry heat indoors in the winter, a requirement for air conditioning during the summer and the effect of spring and fall allergy seasons. As this was a survey-based study, it relied on human recall to evaluate productivity loss, QoL impact and out-of-pocket costs. Recall bias was minimised by using validated recall periods for the VFQ-25 and WPAI questionnaires.^{15 42} Selection bias is a typical limitation of cross-sectional surveys as recruitment of patients with DED who had visited an optometrist/ophthalmologist could lead to an underestimation of productivity losses and their costs (due to patients being too sick to attend a routine visit). An overestimation of losses is also possible because patients with mild DED may not have required a routine visit during the study period. Direct and indirect costs were limited to those pertaining to the patient, and the impact of overall direct costs to the insurer or the overall economy was not considered. Indirect cost according to time lost at work may not account for additional variables such as paid sick leave. The 7-day period over which absenteeism and presenteeism were both estimated and may not accurately reflect patients' actual impairment. The findings of our study are also limited by the sample size and subsequent stratification, particularly for the subgroup with a diagnosis of Sjögren's syndrome ($n=12$). Data related to Sjögren's syndrome should be interpreted with particular caution, as patient-reported absence of this diagnosis was not verified with serology testing or minor salivary gland biopsy. Beyond non-responses to the diagnosis of Sjögren's syndrome, missing or invalid data were minimal, as the investigators checked completed surveys prior to patient discharge. As our analysis was descriptive, and no hypothesis was tested, confounding is not expected to impact our findings.

In conclusion, a cross-sectional survey assessed the economic burden and QoL associated with DED in a sample of patients living in Canada. The annual cost of DED averaged \$24 330, with indirect costs accounting for 90% of the total. Direct and indirect costs, and the negative impact of DED on QoL, were higher in patients with self-reported severe DED compared with those with mild or moderate disease. Sjögren's syndrome was

also associated with higher costs and lower QoL than in patients without Sjögren's syndrome. This study highlights the considerable burdens associated with DED in terms of patient costs (direct and indirect) and reduced QoL, particularly with increased DED severity. These findings in a Canadian population are consistent with those of studies in other countries with different healthcare structures. Further research will be required to establish effective methods of reducing these burdens through expanded use of pharmacological therapies and other interventions such as air quality adjustments or regular work breaks.

Acknowledgements The authors would like to thank the Vector Eye Centre, Calgary, Alberta (Jamie Bhamra, Lead Physician; Alison Endsins, Study Coordinator), Rosedale Medical Centre, Toronto, Ontario (Clara C. Chan, Lead Physician; Paul Lam, Study Coordinator), Ivey Eye Institute, London, Ontario (Rookaya Mather, Lead Physician; Bobbi Smuck and Julie Duncan, Study Coordinators), Clarity Eye Institute, Scarborough, Ontario (King Chow, Lead Physician; Heather Coulter, Study Coordinator), Valley Family Optometry (Krista Flynn, Lead Physician; Shyanne Jenkins, Study Coordinator), and Elmsdale Vision Centre, Elmsdale, Nova Scotia (Andrew Webber, Lead Physician; Dolly Chippett and Stacey MacKenzie, Study Coordinators), and the patients who participated in this study. The authors also wish to thank Johanna Mancini, Megan Naeini and Deepi Minhas of IQVIA for their help in the design, management and analysis of the study, Jeff Alexander of SNELL Medical Communication for writing assistance, and Christopher Reaume of Takeda.

Contributors All authors contributed to the completion of this study and to the preparation of this manuscript.

Funding The study was supported by Takeda and Novartis. The sponsor provided a formal review of the publication; however, the authors had final authority over the content. Medical writing support was provided by Envision Pharma Group, which was supported by Takeda and Novartis.

Competing interests CC has been a consultant for Alcon, Allergan, Bausch+Lomb, Johnson & Johnson, Latician, Santen, and Shire* and has received research support from Allergan, Bausch+Lomb and TearLab. SZ has received grant/research support from Shire* and honoraria/consulting fees from Allergan, Latician, J&J, Santen, and Shire*. VM was a paid consultant to Shire* in connection with this study. JGB was an employee of Shire* at the time of the study and is a current employee of Novartis. CLP has been a consultant for Alcon Canada, Innova Medical, Shire*, Novartis and Bausch+Lomb Canada. *A Takeda company.

Patient consent for publication Not applicable.

Ethics approval The study was performed in accordance with the Health Insurance Portability and Accountability Act 1996 (HIPAA) regulations and the principles of the Declaration of Helsinki. Institutional review board (IRB) and ethics committee approvals were obtained as follows: Rosedale Medical Centre (CCC); Advarra IRB approval SSU00064394 (26 July 2018); Vector Eye Centre (JB); Conjoint Health Research Ethics Board, University of Calgary approval; REB18-1185 (1 August 2018); Ivey Eye Institute (RM); Western University Health Science Research Ethics Board approval 112286 (20 August 2018); Clarity Eye Institute (KC) Advarra IRB approval SSU00075068 (11 January 2019); Valley Family Optometry (KF); Advarra IRB approval SSU00076123 (1 February 2019); Elmsdale Vision Centre (AW); Advarra IRB approval SSU00064416 (26 July 2018); written informed consent was obtained from all patients. Neither patients nor the public were involved in the design, conduct, reporting or dissemination plans of this study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Clara Chan <http://orcid.org/0000-0002-2886-2011>

REFERENCES

- Craig JP, Nichols KK, Akpek EK, *et al.* TFOS DEWS II definition and classification report. *Ocul Surf* 2017;15:276–83.
- Stapleton F, Alves M, Bunya VY, *et al.* TFOS DEWS II epidemiology report. *Ocul Surf* 2017;15:334–65.
- Caffery B, Srinivasan S, Reaume CJ, *et al.* Prevalence of dry eye disease in Ontario, Canada: a population-based survey. *Ocul Surf* 2019;17:526–31.
- Wolffsohn JS, Arita R, Chalmers R, *et al.* TFOS DEWS II diagnostic methodology report. *Ocul Surf* 2017;15:539–74.
- Pflugfelder SC, Solomon A, Stern ME. The diagnosis and management of dry eye: a twenty-five-year review. *Cornea* 2000;19:644–9.
- Jones L, Downie LE, Korb D, *et al.* TFOS DEWS II management and therapy report. *Ocul Surf* 2017;15:575–628.
- Uchino M, Schaumberg DA. Dry eye disease: impact on quality of life and vision. *Curr Ophthalmol Rep* 2013;1:51–7.
- Yu J, Asche CV, Fairchild CJ. The economic burden of dry eye disease in the United States: a decision tree analysis. *Cornea* 2011;30:379–87.
- McDonald M, Patel DA, Keith MS, *et al.* Economic and Humanistic Burden of Dry Eye Disease in Europe, North America, and Asia: A Systematic Literature Review. *Ocul Surf* 2016;14:144–67.
- Abetz L, Rajagopalan K, Mertzanis P, *et al.* Development and validation of the impact of dry eye on everyday life (IDEEL) questionnaire, a patient-reported outcomes (pro) measure for the assessment of the burden of dry eye on patients. *Health Qual Life Outcomes* 2011;9:111.
- Schaumberg DA, Gulati A, Mathers WD, *et al.* Development and validation of a short global dry eye symptom index. *Ocul Surf* 2007;5:50–7.
- Reilly Associates. WPAI:SHP v2.0 (US English), 2010. Available: http://www.reillyassociates.net/WPAI_SHP.html
- Statistics Canada. 2016 Census of Population - Employment Income Statistics (table) [Internet], 2017. Available: <http://www12.statcan.gc.ca/census-recensement/2016/dp-pd/td-t/Rp-eng.cfm?LANG=E&APATH=3&DETAIL=0&DIM=0&FL=A&FREE=0&GC=0&GID=0&GK=0&GRP=1&PID=110647&PRID=10&PTYPE=109445&S=0&SHOWALL=0&SUB=0&Temporal=2017&THEME=123&VID=0&VNAMEE=&VNAMEF>
- Bank of Canada. Consumer Price Indexes for Canada [Internet]. Available: <https://www.bankofcanada.ca/rates/related/inflation-calculator/>
- Reilly MC, Zbrozek AS, Dukes EM. The validity and reproducibility of a work productivity and activity impairment instrument. *Pharmacoeconomics* 1993;4:353–65.
- Reilly MC, Bracco A, Ricci J-F, *et al.* The validity and accuracy of the Work Productivity and Activity Impairment questionnaire--irritable bowel syndrome version (WPAI:IBS). *Aliment Pharmacol Ther* 2004;20:459–67.
- Reilly MC, Gooch KL, Wong RL, *et al.* Validity, reliability and responsiveness of the work productivity and activity impairment questionnaire in ankylosing spondylitis. *Rheumatology* 2010;49:812–9.
- Reilly MC, Gerlier L, Brabant Y, *et al.* Validity, reliability, and responsiveness of the work productivity and activity impairment questionnaire in Crohn's disease. *Clin Ther* 2008;30:393–404.
- Revicki DA, Willian MK, Menter A, *et al.* Impact of adalimumab treatment on patient-reported outcomes: results from a phase III clinical trial in patients with moderate to severe plaque psoriasis. *J Dermatol Treat* 2007;18:341–50.
- Chen H, Blanc PD, Hayden ML, *et al.* Assessing productivity loss and activity impairment in severe or difficult-to-treat asthma. *Value Health* 2008;11:231–9.
- Zhang W, Bansback N, Boonen A, *et al.* Validity of the work productivity and activity impairment questionnaire--general health version in patients with rheumatoid arthritis. *Arthritis Res Ther* 2010;12:R177.
- Nichols KK, Bacharach J, Holland E, *et al.* Impact of dry eye disease on work productivity, and patients' satisfaction with over-the-counter dry eye treatments. *Invest Ophthalmol Vis Sci* 2016;57:2975–82.
- Dana R, Meunier J, Markowitz JT, *et al.* Patient-Reported burden of dry eye disease in the United States: results of an online cross-sectional survey. *Am J Ophthalmol* 2020;216:7–17.
- Patel VD, Watanabe JH, Strauss JA, *et al.* Work productivity loss in patients with dry eye disease: an online survey. *Curr Med Res Opin* 2011;27:1041–8.
- RAND. The National eye Institute 25-Item visual function questionnaire (VFQ-25) manual, 2000. Available: https://www.rand.org/content/dam/rand/www/external/health/surveys_tools/vfq/vfq25_manual.pdf
- Farrand KF, Stillman IÖ, Fridman M, Schaumberg S, *et al.* Impact of dry eye disease on quality of life, work productivity, daily activities, and health care resource use in a survey of 74,095 American adults. *Value Health* 2016;19:A127.
- Gardner BT, Dale AM, Buckner-Petty S, *et al.* Comparison of employer productivity metrics to lost productivity estimated by commonly used questionnaires. *J Occup Environ Med* 2016;58:170–7.
- Zhang W, Gignac MAM AM, Beaton D, *et al.* Canadian arthritis network work productivity group. Productivity loss due to presenteeism among patients with arthritis: estimates from 4 instruments. *J Rheumatol* 2010;37:1805–14.
- Armstrong AW, Lynde CW, McBride SR, *et al.* Effect of ixekizumab treatment on work productivity for patients with moderate-to-severe plaque psoriasis: analysis of results from 3 randomized phase 3 clinical trials. *JAMA Dermatol* 2016;152:661–9.
- Clegg JP, Guest JF, Lehman A, *et al.* The annual cost of dry eye syndrome in France, Germany, Italy, Spain, Sweden and the United Kingdom among patients managed by ophthalmologists. *Ophthalmic Epidemiol* 2006;13:263–74.
- Online currency converter. Available: <https://freecurrencyrates.com/en/exchange-rate-history/USD-CAD/2006/cbr> [Accessed 29 Apr 2021].
- Cornec D, Devauchelle-Pensec V, Mariette X, *et al.* Severe health-related quality of life impairment in active primary Sjögren's syndrome and patient-reported outcomes: data from a large therapeutic trial. *Arthritis Care Res* 2017;69:528–35.
- Nichols KK, Mitchell GL, Zadnik K. Performance and repeatability of the NEI-VFQ-25 in patients with dry eye. *Cornea* 2002;21:578–83.
- Mertzanis P, Abetz L, Rajagopalan K, *et al.* The relative burden of dry eye in patients' lives: comparisons to a U.S. normative sample. *Invest Ophthalmol Vis Sci* 2005;46:46–50.
- Tellefsen Nøland S, Badian RA, Utheim TP, *et al.* Sex and age differences in symptoms and signs of dry eye disease in a Norwegian cohort of patients. *Ocul Surf* 2021;19:68–73.
- Garcia-Alfaro P, Garcia S, Rodriguez I, *et al.* Dry eye disease symptoms and quality of life in perimenopausal and postmenopausal women. *Climacteric* 2021;24:261–6.
- Wang MTM, Muntz A, Mamidi B, *et al.* Modifiable lifestyle risk factors for dry eye disease. *Cont Lens Anterior Eye* 2021;101409.
- Sánchez-Valerio MDR, Mohamed-Noriega K, Zamora-Ginez I, *et al.* Dry eye disease association with computer exposure time among subjects with computer vision syndrome. *Clin Ophthalmol* 2020;14:4311–7.
- Uchino M, Schaumberg DA, Dogru M, *et al.* Prevalence of dry eye disease among Japanese visual display terminal users. *Ophthalmology* 2008;115:1982–8.
- Gajta A, Turkoanje D, Malaescu I. Dry eye syndrome among computer users. *AIP Conference Proceedings* 2015;1694:040011.
- Moon JH, Kim KW, Moon NJ. Smartphone use is a risk factor for pediatric dry eye disease according to region and age: a case control study. *BMC Ophthalmol* 2016;16:188.
- Mangione CM, Lee PP, Pitts J, *et al.* Psychometric properties of the National eye Institute visual function questionnaire (NEI-VFQ). NEI-VFQ field test Investigators. *Arch Ophthalmol* 1998;116:1496–504.

SECTION A: GENERAL QUESTIONS

A1. Today's date

Day		Month		Year			

A2. What is your birth month and birth year?

Month		Year			

A3. What is your sex?

- Male
 Female

A4. What is your ethnicity? You can put an in more than one box, if applicable.

- Caucasian
 Latino/Hispanic
 Middle Eastern
 African
 Caribbean
 South Asian
 East Asian
 Other _____

A5. What is the highest level of education you have achieved?

Look for your highest degree in education and fill in an x in the box.

- No certificate, diploma or degree
 High school diploma or equivalency certificate
 Trades: Certificate or diploma (excluding certificate of apprenticeship)
 Trades: Certificate of Apprenticeship or Certificate of Qualification
 College, CEGEP or other non-university certificate or diploma
 University: Certificate or diploma BELOW bachelor level
 University: Certificate, diploma or degree AT bachelor level
 University: Certificate, diploma or degree ABOVE bachelor level

A6. What is your employment status? Place an in the box for what you usually do.

- I am employed
- I am unemployed – looking for a job
- I am unemployed – not looking for a job
- I am on disability leave from work due to Dry Eye Disease (DED)
- I am on disability leave from work due to other reasons
- I am retired or on a pre-pension plan
- I do something else, namely: _____

SECTION B: HEALTH QUESTIONS

B1. Do you smoke?

- Previous smoker, but stopped
- Yes – current smoker
- No – never smoked

B2. Are you currently pregnant?

- Yes
- No

B3. Do you wear contact lenses?

- Yes
- No

B4. How many hours per day do you estimate you spend looking at a screen?

(TV, phone, iPad/tablet, computer, etc.)

- 0 hours
- Less than 1 hour, more than 0
- 1-2 hours
- 3-4 hours
- 5-6 hours
- Other: ____ hours

B5. Which type of treatment are you currently applying to your eyes?

Select all that apply.

- Preserved
- Non-preserved (ex. Lacrisert)
- Prescription eye drops (ex. Xiidra, Restasis)
- Specialty drugs
- Compounded ointments
- Gel (ex. Lotemax)
- Other _____
- I don't know
- None

B6. What other medications are you currently taking? Select all that apply.

- | | |
|---|---|
| <input type="checkbox"/> Anti-depressants | <input type="checkbox"/> Beta blockers |
| <input type="checkbox"/> Anti-Cholinergics (e.g. Parkinson's medications, antipsychotics, sleeping pills) | <input type="checkbox"/> Oral contraceptives |
| <input type="checkbox"/> Antihistamines | <input type="checkbox"/> Diuretics |
| <input type="checkbox"/> Anti-anxiety | <input type="checkbox"/> Chemotherapy/radiation therapy |
| <input type="checkbox"/> Botox | <input type="checkbox"/> Other _____ |
| <input type="checkbox"/> Hormone replacement therapy | <input type="checkbox"/> I don't know |
| <input type="checkbox"/> Isotretinoin (e.g. Accutane) | <input type="checkbox"/> None |
| <input type="checkbox"/> Multivitamins | |

B7. Have you been formally diagnosed with Sjögren syndrome?

- Yes
 No
 I don't know

B8. Do you currently have any of the following conditions? Select all that apply.

- | | |
|--|--|
| <input type="checkbox"/> Androgen deficiency | <input type="checkbox"/> Mental health conditions (<i>i.e.</i> mood, depression, anxiety) |
| <input type="checkbox"/> Autoimmune diseases (<i>e.g.</i> Sjögren, RA, lupus) | <input type="checkbox"/> Migraines |
| <input type="checkbox"/> Chronic pain | <input type="checkbox"/> Rosacea |
| <input type="checkbox"/> Diabetes | <input type="checkbox"/> Sick Building Syndrome |
| <input type="checkbox"/> Fibromyalgia | <input type="checkbox"/> Thyroid disorder |
| <input type="checkbox"/> Gout | <input type="checkbox"/> Vitamin A deficiency |
| <input type="checkbox"/> Hematopoietic stem cell transplantation or Graft vs. Host Disease | <input type="checkbox"/> Other _____ |
| <input type="checkbox"/> HIV | <input type="checkbox"/> None |
| <input type="checkbox"/> Irritable Bowel Syndrome | |

B9. How long have you been suffering from Dry Eye Disease?

- For less than 1 year
 For the past 1-5 years
 For the past 6-10 years
 Other _____ years

B10. Other than Dry Eye Disease, do you have any of the following conditions?

Select all that apply.

- Glaucoma
- Keratoconus +/- corneal collagen crosslinking
- Meibomian gland dysfunction
- Prior conjunctivitis (*pink eye*)
- Seasonal allergies with itchy eyes
- Surfer's Eye (*pterygium*)
- Other _____
- None

B11. Have you had any ocular surgeries or injections?

- Glaucoma surgery
- Cataract
- Eye lid surgery
- Refractive eye surgery, including LASIK or PRK
- Other _____
- None

SECTION C: EYE DRYNESS SCORE

Visual Analogue Scale (VAS)

Please rate your discomfort with the *Eye Dryness* symptom on a scale from 0 to 100 by placing a vertical mark (|) on the horizontal line indicating your level of eye discomfort

0 corresponds to: “no discomfort”

100 corresponds to: “maximal (the most) discomfort”

Eye Dryness



SECTION D: VISUAL FUNCTIONING QUESTIONNAIRE – 25

The following is a survey with statements about problems which involve your vision or feelings that you have about your vision condition. After each question please choose the response that best describes your situation.

Please answer all the questions as if you were wearing your glasses or contact lenses (if any).

Please take as much time as you need to answer each question. All your answers are confidential. In order for this survey to improve our knowledge about vision problems and how they affect your quality of life, your answers must be as accurate as possible. Remember, if you wear glasses or contact lenses, please answer all of the following questions as though you were wearing them.

PART 1 – GENERAL HEALTH AND VISION

1. In general, would you say your overall health is:

(Circle One)

Excellent

Very Good.....

Good.....

Fair

Poor.....

2. At the present time, would you say your eyesight using both eyes (with glasses or contact lenses, if you wear them) is excellent, good, fair, poor, or very poor or are you completely blind?

(Circle One)

Excellent

Good.....

Fair

Poor.....

Very Poor

Completely Blind.....

3. How much of the time do you worry about your eyesight?

(Circle One)

- None of the time 1
 A little of the time 2
 Some of the time 3
 Most of the time 4
 All of the time? 5

4. How much pain or discomfort have you had in and around your eyes (for example, burning, itching, or aching)? Would you say it is:

(Circle One)

- None 1
 Mild 2
 Moderate 3
 Severe, or 4
 Very severe? 5

PART 2 – DIFFICULTY WITH ACTIVITIES

The next questions are about how much difficulty, if any, you have doing certain activities wearing your glasses or contact lenses if you use them for that activity.

5. How much difficulty do you have reading ordinary print in newspapers? Would you say you have:

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

6. How much difficulty do you have doing work or hobbies that require you to see well up close, such as cooking, sewing, fixing things around the house, or using hand tools?
Would you say:

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

7. Because of your eyesight, how much difficulty do you have finding something on a crowded shelf?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

8. How much difficulty do you have reading street signs or the names of stores?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

9. Because of your eyesight, how much difficulty do you have going down steps, stairs, or curbs in dim light or at night?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

10. Because of your eyesight, how much difficulty do you have noticing objects off to the side while you are walking along?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

11. Because of your eyesight, how much difficulty do you have seeing how people react to things you say?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

12. Because of your eyesight, how much difficulty do you have picking out and matching your own clothes?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

13. Because of your eyesight, how much difficulty do you have visiting with people in their homes, at parties, or in restaurants?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

14. Because of your eyesight, how much difficulty do you have going out to see movies, plays, or sports events?

(Circle One)

- No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4
 Stopped doing this because of your eyesight..... 5
 Stopped doing this for other reasons or not interested in doing this 6

15. Are you currently driving, at least once in a while?*(Circle One)*

Yes 1 *Skip To Q 15c*
 No 2

15a. IF NO: Have you never driven a car or have you given up driving?*(Circle One)*

Never drove 1 *Skip To Part 3, Q 17*
 Gave up..... 2

15b. IF YOU GAVE UP DRIVING: Was that mainly because of your eyesight, mainly for some other reason, or because of both your eyesight and other reasons?*(Circle One)*

Mainly eyesight 1 *Skip To Part 3, Q 17*
 Mainly other reasons 2 *Skip To Part 3, Q 17*
 Both eyesight and other reasons 3 *Skip To Part 3, Q 17*

15c. IF CURRENTLY DRIVING: How much difficulty do you have driving during the daytime in familiar places? Would you say you have:*(Circle One)*

No difficulty at all 1
 A little difficulty 2
 Moderate difficulty 3
 Extreme difficulty 4

16. How much difficulty do you have driving at night? Would you say you have:

(Circle One)

- No difficulty at all 1
- A little difficulty 2
- Moderate difficulty 3
- Extreme difficulty 4
- Have you stopped doing this because of your eyesight 5
- Have you stopped doing this for other reasons or are you not interested in doing this 6

16A. How much difficulty do you have driving in difficult conditions, such as in bad weather, during rush hour, on the freeway, or in city traffic? Would you say you have:

(Circle One)

- No difficulty at all 1
- A little difficulty 2
- Moderate difficulty 3
- Extreme difficulty 4
- Have you stopped doing this because of your eyesight 5
- Have you stopped doing this for other reasons or are you not interested in doing this 6

PART 3: RESPONSES TO VISION PROBLEMS

The next questions are about how things you do may be affected by your vision. For each one, please circle the number to indicate whether for you the statement is true for you all, most, some, a little, or none of the time.

(Circle One On Each Line)

READ CATEGORIES:	All of the time	Most of the time	Some of the time	A little of the time	None of the time
17. <u>Do you accomplish less than you would like because of your vision?</u>	1	2	3	4	5
18. <u>Are you limited in how long you can work or do other activities because of your vision?</u>	1	2	3	4	5
19. <u>How much does pain or discomfort in or around your eyes, for example, burning, itching, or aching, keep you from doing what you'd like to be doing? Would you say:</u>	1	2	3	4	5

For each of the following statements, please circle the number to indicate whether for you the statement is definitely true, mostly true, mostly false, or definitely false for you or you are not sure.

	(Circle One On Each Line)				
	Definitely True	Mostly True	Not Sure	Mostly False	Definitely False
20. I <u>stay home most of the time</u> because of my eyesight	1	2	3	4	5
21. I feel <u>frustrated</u> a lot of the time because of my eyesight	1	2	3	4	5
22. I have <u>much less control</u> over what I do, because of my eyesight	1	2	3	4	5
23. Because of my eyesight, I have to <u>rely too much on what other people tell me</u>	1	2	3	4	5
24. I <u>need a lot of help</u> from others because of my eyesight	1	2	3	4	5
25. I worry about doing things that <u>will embarrass myself or others</u> , because of my eyesight	1	2	3	4	5

SECTION E: WORK PRODUCTIVITY AND ACTIVITY IMPAIRMENT QUESTIONNAIRE

The following questions ask about the effect of your **Dry Eye Disease (DED)** on your ability to work and perform regular activities. *Please fill in the blanks or circle a number, as indicated.*

1. Are you currently employed (working for pay)? _____ NO ___ YES
If NO, check "NO" and skip to question 6.

The next questions are about the **past seven days**, not including today.

2. During the past seven days, how many hours did you miss from work because of problems associated with your DED? Include hours you missed on sick days, times you went in late, left early, etc., because of your DED. Do not include time you missed to participate in this study.

_____ HOURS

3. During the past seven days, how many hours did you miss from work because of any other reason, such as vacation, holidays, time off to participate in this study?

_____ HOURS

4. During the past seven days, how many hours did you actually work?

_____ HOURS (If "0", skip to question 6.)

5. During the past seven days, how much did your DED affect your productivity while you were working?

Think about days you were limited in the amount or kind of work you could do, days you accomplished less than you would like, or days you could not do your work as carefully as usual. If DED affected your work only a little, choose a low number. Choose a high number if DED affected your work a great deal.

Consider only how much DED affected
productivity while you were working.

DED had no effect
on my work

0 1 2 3 4 5 6 7 8 9 10

DED completely
prevented me from
working

CIRCLE A NUMBER

6. During the past seven days, how much did your DED affect your ability to do your regular daily activities, other than work at a job?

By regular activities, we mean the usual activities you do, such as work around the house, shopping, childcare, exercising, studying, etc. Think about times you were limited in the amount or kind of activities you could do and times you accomplished less than you would like. If DED affected your activities only a little, choose a low number. Choose a high number if DED affected your activities a great deal.

Consider only how much DED affected your ability to do your regular daily activities, other than work at a job.



CIRCLE A NUMBER

SECTION F: OUT-OF-POCKET EXPENSES

We want to better understand the financial burden that your Dry Eye Disease is having on you. The following questions ask about the amount of money you spend out of your own pocket when buying treatments such as eye drops or supplements, and any extra money you spend to see your ophthalmologist/optometrist.

RESOURCE A: Over-the-Counter Ocular Lubricants - Artificial tears

(Examples: Systane, Refresh, Blink, Bion, Hydrasense, Hylo)

1. Have you used NON-PRESCRIPTION ARTIFICIAL TEARS in the past 3 (THREE) months?

- Yes
- No – skip to *RESOURCE B, Question #5*

2. If so, how often do you usually use NON-PRESCRIPTION ARTIFICIAL TEARS daily?

- 1-2 times a day
- 3-4 times a day
- 5-6 times a day
- Other: ____ times a day
- I don't know

3. Are you paying for NON-PRESCRIPTION ARTIFICIAL TEARS out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to *RESOURCE B, Question #5*
- I don't know

4. If so, how much money PER ONE MONTH are you typically spending on NON-PRESCRIPTION ARTIFICIAL TEARS?

- \$10-19 per month
- \$20-29 per month
- \$30-39 per month
- \$40-49 per month
- Other: \$ ____ per month
- I don't know

RESOURCE B: Over-the-Counter Ocular Lubricants - Gels or ointments*(Example: Teargel, Duolube, Lacrilube, Ocunox)***5. Have you used NON-PRESCRIPTION OCULAR GELS/OINTMENTS in the past 3 (THREE) months?**

- Yes
- No – skip to RESOURCE C, Question #9

6. If so, how often do you usually use NON-PRESCRIPTION OCULAR GELS/OINTMENTS daily?

- 1-2 times a day
- 3-4 times a day
- 5-6 times a day
- Other: ____ times a day
- I don't know

7. Are you paying for NON-PRESCRIPTION OCULAR GELS/OINTMENTS out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to RESOURCE C, Question #9
- I don't know

8. If so, how much money PER ONE MONTH are you typically spending on NON-PRESCRIPTION OCULAR GELS/OINTMENTS?

- \$10-19 per month
- \$20-29 per month
- \$30-39 per month
- \$40-49 per month
- \$50-59 per month
- Other: \$ ____ per month
- I don't know

RESOURCE C: Prescription Ocular Lubricants

(Example: serum tears, amniotic fluid extract (Regenereyes), albumin drops (specifically from princess Margaret hospital).

9. Have you used PRESCRIPTION OCULAR LUBRICANT in the past 3 (THREE) months?

- Yes
- No – skip to RESOURCE D, Question #13

10. If so, how often do you usually use PRESCRIPTION OCULAR LUBRICANT daily?

- 1-2 times a day
- 3-4 times a day
- 5-6 times a day
- Other: ____ times a day
- I don't know

11. Are you paying for PRESCRIPTION OCULAR LUBRICANT out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to RESOURCE D, Question #13
- I don't know

12. If so, how much money PER ONE MONTH are you typically spending on PRESCRIPTION OCULAR LUBRICANT?

- \$40-49
- \$59-149
- \$150-169
- Other: \$____ per month
- I don't know

RESOURCE D: Prescription Ocular Anti-inflammatory Agents*(Example: Xiidra, Restasis, Cyclosporine, steroids, lotemax)***13. Have you used PRESCRIPTION OCULAR ANTI-INFLAMMATORY AGENTS in the past 3 (THREE) months?**

- Yes
- No – skip to RESOURCE E, Question #17

14. If so, how often do you usually use PRESCRIPTION OCULAR ANTI-INFLAMMATORY AGENTS daily?

- 1-2 times a day
- 3-4 times a day
- 5-6 times a day
- Other: ____ times a day
- I don't know

15. Are you paying for PRESCRIPTION OCULAR ANTI-INFLAMMATORY AGENTS out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to RESOURCE E, Question #17
- I don't know

16. If so, how much money PER ONE MONTH are you typically spending on PRESCRIPTION OCULAR ANTI-INFLAMMATORY AGENTS?

- \$0-\$49 per month
- \$50-\$99 per month
- \$100-\$149 per month
- \$150-\$199 per month
- \$200-\$249 per month
- Other: \$ ____ per month
- I don't know

RESOURCE E: Nutritional Supplements*(Example: fish oil, flax seed oil)*

17. Have you used this resource NUTRITIONAL SUPPLEMENTS TO IMPROVE DRY EYE in the past 3 (THREE) months?

- Yes
- No – skip to *RESOURCE F, Question #21*

18. If so, how often do you usually use NUTRITIONAL SUPPLEMENTS TO IMPROVE DRY EYE?

- Everyday
- Every other day
- Twice a week
- Once a week
- Once a month
- Other: ___ times per _____
- I don't know

19. Are you paying for NUTRITIONAL SUPPLEMENTS TO IMPROVE DRY EYE out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to *RESOURCE F, Question #21*
- I don't know

20. If so, how much money PER ONE MONTH are you typically spending on NUTRITIONAL SUPPLEMENTS TO IMPROVE DRY EYE?

- \$0-\$9 per month
- \$10-\$19 per month
- \$20-\$29 per month
- \$30-\$39 per month
- \$40-\$49 per month
- Other: \$___ per month
- I don't know

RESOURCE F: Punctal Plugs**21. Have you had PUNCTAL PLUGS inserted in one or both of your eyes before?**

- Yes
- No – skip to *RESOURCE G, Question #25*

22. If so, how many PUNCTAL PLUG implants have you had inserted?

(For example, if you have received 1 implant in each eye, the total number of implants you've receive is 2. If one of those implants fell out and you had to get it replaced, then the total number of implants you've ever had equals 3)

- Only 1 implant in my life
- 2 implants in total in my life
- 3 implants in total in my life
- 4 implants in total in my life
- Other: ___ implants
- I don't know

23. Did you pay for your PUNCTAL PLUG IMPLANTS out-of-pocket?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to *RESOURCE G, Question #25*
- I don't know

24. If so, how much money, on average, did you spend on PUNCTAL PLUG IMPLANTS?

- \$0-\$49 per implant
- \$50-\$99 per implant
- \$100-\$149 per implant
- \$150-\$199 per implant
- \$200-\$249 per implant
- Other \$_____ per implant
- I don't know

RESOURCE G: Optometrist Visits**25. Is TODAY your FIRST OPTOMETRIST VISIT for dry eye disease?**

- Yes – skip to *RESOURCE H Question #29*
- No

26. If NOT, how many times did you VISIT THE OPTOMETRIST in the last 2 (TWO) YEARS for dry eye disease?

- Twice a year
- Once a year
- Once every two years
- Other: ___ times per ___
- I don't know

27. Do you pay anything out-of-pocket during these visits to the OPTOMETRIST?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – skip to *RESOURCE H, Question #29*
- I don't know

28. If so, how much money on average PER VISIT?

- \$0-\$49 per visit
- \$50-\$99 per visit
- \$100-\$149 per visit
- \$150-\$199 per visit
- \$200-\$249 per visit
- Other \$___ per visit
- I don't know

RESOURCE H: Ophthalmologist Visits**29. Is TODAY your FIRST OPHTHALMOLOGIST visit for dry eye disease?**

- Yes – *skip to the end of the survey*
- No

30. If NOT, how often did you VISIT THE OPHTHALMOLOGIST in the last 2 (TWO) YEARS for dry eye disease?

- Twice a year
- Once a year
- Once every two years
- Other: ___ times per ___
- I don't know

31. Do you pay anything out-of-pocket to VISIT THE OPHTHALMOLOGIST?

- Yes, because I either don't have insurance or my insurance does not cover this
- Partially, since insurance covers some costs
- No, insurance pays for everything – *skip to the end of the survey*
- I don't know

32. If so, how much money on average PER VISIT?

- \$0
- \$1-\$49 per visit
- \$50-\$99 per visit
- \$100-\$149 per visit
- \$150-\$199 per visit
- \$200-\$249 per visit
- Other \$___ per visit
- I don't know

That was the last question.**What should you do with the survey now that it is filled in?**

Please return the completed survey and return it to your optometrist/ophthalmologist. Please note that all your answers will be treated as strictly confidential.

Thank you very much!

Table S2. Indirect cost outcomes from the Work Productivity and Activity Impairment survey¹²

Absenteeism			
Percent work time missed due to problem, %	Continuous	E2, E4	$E2/(E2+E4)$
Average annual cost of missed work time, \$/year (SD)	Continuous	E2, E4, Statistics Canada ¹³	Variable 1 multiplied by the patient's annual salary, inferred from education level, age, and sex ¹³
Presenteeism			
Percent impairment while working due to problem	Continuous	E5	$E5/10$
Average annual cost due to decrease in work productivity (\$/year, SD)	Continuous	E5, Statistics Canada ¹³	Variable #3 multiplied by the patient's annual salary, inferred from education level, age, and sex ¹³
Percent activity impairment due to problem	Continuous	E6	$E6/10$
Percent overall work impairment due to problem	Continuous	E2, E4, E5	$E2/(E2+E4) + [(1 - (E2/(E2+E4))) \times (E5/10)]$

SD, standard deviation