

# P18-A148 ASSYMETRICAL INJECTOR FOR ENDOTHELIUM-IN DMEK WITHOUT THE NEED OF PULL-THROUGH TECHNIQUE

<sup>1,2</sup>Mikhail Tsurkan, <sup>3</sup>John Lohmeier, <sup>3</sup>Staci Terrin, <sup>2</sup>Simone Arndt, <sup>2</sup>Sarah Tsurkan. <sup>1</sup>TissueGUARD GmbH, Dresden, Germany; <sup>2</sup>Leibniz-Institut für Polymerforschung, Dresden, Germany; <sup>3</sup>Rocky Mountain Lions Eye Bank, Aurora, USA

10.1136/bmjophth-2023-EEBA.17

**Purpose** One of latest surgical development of preloaded Descemet membrane endothelial keratoplasty (DMEK) is the delivery of the graft with the endothelium inwards, which allows for a very fast operation, but requires a pull-through surgical technique. Although the tri-folded, endo-in DMEK technique has significant advantages, the absence of proper surgical instruments that could allow their use without the 'pull-through' technique still restricts the wide use of such an operation. None of the available commercial DMEK injectors could be used for tri-folded DMEK (endothelium-inward) orientation, as it requires the graft to be intently secured within the injector. This report presents a retrospective eye bank validation study of an asymmetrical injector designed to orientally implant a tri-folded DMEK graft without needing a pull-through technique.

**Methods** The injector is made from transparent plastic, allowing microscopic tissue validation directly before injection. The device is asymmetrical, so the orientation of the graft can be controlled and validated according to the best eye bank practice, which is critical for successful tri-folded DMEK graft clinical application. Four different designs of the internal compartment of the injectors were evaluated with DMEK tissues. Mates from two pairs were tested on each device type, totaling 16 grafts, all loaded with folded, endo-in grafts. The tissue was prepared, loaded into the injector, and ejected to imitate the tissue manipulation in DMEK operation.

**Results** After graft loading the delivery of the endothelium-in grafts was performed by injection, without the need for a pull-through technique. One graft (6.25%) has double-scrolled (changed its folding) within the injector with a larger (1.5 mm) internal compartment. The loss of valuable cells was between 3-23% (13.98% average). No significant differences in cell loss were observed between injectors with different internal compartment sizes. Higher viability loss (17.3% +/- 5.7) was observed for the grafts with >20 days death to prep-days in comparison with grafts stored with less than two weeks (10.9% +/- 2.1).

**Conclusion** The TissueGUARD injector is the only injector that currently allows oriented, tri-folded DMEK injection without the need for a pull-through technique. The average cell loss after loading and ejection was 13.98%, which is comparable/better than the current best practice with the pre-cut-preloaded technique of naturally folded DMEK.

# P19-A118 BOWMAN LAYER ONLAY: PREPARATION AND TRANSPLANTATION (BLOT)

<sup>1,2</sup>Esther A Groeneveld-van Beek, <sup>1,3</sup>Lydia van der Star, <sup>1,2,3</sup>Paulina Bylewska, <sup>1</sup>Maloeke de Jong, <sup>1</sup>Silke Oellerich, <sup>1,2</sup>Jacqueline van der Wees, <sup>1,3</sup>Isabel Dapena, <sup>1,2,3</sup>Gerrit RJ Melles, <sup>1,2,3,4</sup>Viridiana Kocaba. <sup>1</sup>Netherlands Institute for Innovative Ocular Surgery, Rotterdam, The Netherlands; <sup>2</sup>Amnitrans EyeBank Rotterdam, The Netherlands; <sup>3</sup>Melles Cornea Clinic Rotterdam, The Netherlands; <sup>4</sup>Tissue and Cell Therapy Group, Singapore Eye Research Institute, Singapore

10.1136/bmjophth-2023-EEBA.18

**Purpose** With the introduction of Bowman layer onlay transplantation (BLOT), the need for BL transplants increases.

In this study, the clinical outcomes of BLOT are described and the results of three different BL graft preparation methods are evaluated: manually (m-BL), femtosecond laser-assisted (fs-BL), and femtosecond laser-assisted followed by excimer laser (fs/ex-BL).

**Method** Twenty-one eyes with advanced progressive keratoconus underwent BLOT with m-BL. Best spectacle- and/or best contact lens-corrected visual acuity (BSCVA/BCLVA), corneal tomography, and complications were recorded. Follow-up ranged from 6-36 months with a mean follow-up time of 21 ± 12 months.

To evaluate BL preparation methods, Descemet membrane-denuded donor corneas (n=41) were used (n=2 for m-BL, n=18 for fs-BL and n=21 for fs/ex-BL). For fs-BL, corneas were placed on an artificial anterior chamber and different depth cuts were performed with decreasing decrements starting from 30 µm (diameter 9.0 mm). For fs/ex-BL, a superficial flap of 80 µm was created by the femtosecond laser (FEMTO-LDV Z8, Ziemer). Followed by residual stroma ablation by excimer laser (Schwind Amaris 750S) with increasing increments. Grafts were analyzed visually, and graft thickness regularity was evaluated by histological analysis and Transmission Electron Microscopy (TEM).

**Results** All twenty-one surgeries could be performed without intraoperative complications. Average maximum keratometry changed from 75.8±12D preoperatively to 72.2±9D at the last available follow-up (n=21, P<0.05), and BSCVA/BCLVA improved. Five patients required a re-graft; four of those because of a graft detachment within one week.

Evaluation of BL-preparation methods: Fs-BL preparation was successful until 14µm cuts (success rate: 12 out of 14, 86%). Fs/ex-BL graft preparation was most successful after an 80µm cut by femtosecond laser with subsequent 60µm ablation by excimer laser (success rate: 15 out of 21, 71%). After the femtosecond laser cut, traces of the femtosecond laser treatment were visible on the flap. While m-BL showed long protruding stromal fibers, they were shorter in fs-BL and absent in fs/ex-BL.

**Conclusion** BL-onlay grafting may be a feasible surgical technique, providing on average -3D of corneal flattening in eyes with advanced progressive keratoconus, while improving patient's visual acuity.

Fs-BL and fs/ex-BL preparation may be faster alternatives to manual BL graft preparation.

# P20-A129 AN INNOVATIVE WOUND HEALING METHOD REVEALS A DONOR AND POST-MORTEM TIME-DEPENDENT REGENERATION OF CORNEAL EPITHELIUM

<sup>1</sup>Filippo Bonelli, <sup>1,2</sup>Umberto Rodella, <sup>1</sup>Elisa Fasolo, <sup>1</sup>Vanessa Barbaro, <sup>1</sup>Ilaria Zorzi, <sup>2</sup>Jana D'Amato Tóthová, <sup>1</sup>Stefano Ferrari. <sup>1</sup>Fondazione Banca degli Occhi del Veneto (FBOV), Italy; <sup>2</sup>Research and Development, AL.CHI.M.I.A. S.R.L, Italy

10.1136/bmjophth-2023-EEBA.19

**Purpose** The aim of this study was to establish and optimize a new and reproducible epithelial wound healing model on human corneas. This assay was used to study the kinetics of epithelial regeneration following a chemical injury.

**Methods** Thirty (n=30) human corneas unsuitable for transplant were used for the experiments. Corneas were cultured in Storagix medium (FBOV) at 31°C. Epithelial integrity