

	FASD Eye Code Protocol	
Name		
Code score	A. Best corrected visual acuity (BCVA)	Patient score
1	BCVA worst eye logMAR ≤ 0.1 (decimal ≥ 0.8)	
2	BCVA best eye logMAR ≤ 0.5 (decimal ≥ 0.65)	
3	BCVA each eye/best eye logMAR ≤ 0.5 and ≥ 0.3 (decimal ≥ 0.3 and ≤ 0.5)	
4	BCVA best eye logMAR > 0.5 (decimal < 0.3)	
	B. Refraction in cycloplegia	
1	No significant refractive error	
2	Hyperopia (≥ 2.0 D SE) or myopia (≥ 1.0 D SE) in one or both eyes	
3	Anisometropia (≥ 1.0 D SE)	
4	Astigmatism (> 1.0 D) in one or both eyes	
	C. Strabismus and binocular function	
1	Orthophoria or heterophoria and normal stereo acuity (TNO $\leq 60''$; Lang $\leq 200''$)	
2	Heterophoria or intermittent heterotropia and normal stereo acuity (TNO $\leq 60''$; Lang $\leq 200''$)	
3	No apparent deviation, or heterophoria, or intermittent heterotropia and subnormal stereo acuity (TNO $> 60''$; Lang $> 200''$)	
4	Heterotropia and subnormal stereo acuity (TNO $> 60''$; Lang $> 200''$) or no stereo acuity (TNO neg; Lang neg)	
	D. Ocular structural abnormalities	
1	No ocular structural abnormalities	
2	Ptosis and/or epicanthic folds in one or both eyes	
3	Ocular abnormalities of the retinal vessels (i.e. increased tortuosity) and/or disc (excluding optic nerve hypoplasia)	
4	Optic nerve hypoplasia (ONH in one or both eyes)	
	Total score (A+B+C+D)	