

Supplemental Table 2: Change in prevalence of optical coherence tomography (OCT) features in patients with Best vitelliform macular dystrophy over time.

	Baseline	Follow-up	p-value
Number of Eyes	65		-
Follow-up Period [Years] (mean, median, range)	5.1, 5, 1 to 11		-
Age [Years] (mean, median, range)	40.1, 42.0, 4 to 84	45.1, 50.0, 8 to 89	0.12
LogMAR BCVA (mean, median, range)	0.34, 0.3, -0.12 to 1.0	0.27, 0.20, -0.52 to 0.90	0.18
% Eyes with BCVA \geq 20/50	60	65	0.33
CST [μm] (mean, median, range)	359.7, 329, 177 to 627	340, 321, 142 to 629	0.17
<u>Feature Rate (%)</u>			
Pre-VL	10.8	3.1	0.23
Solid VL	10.8	15.4	0.33
SRF	7.7	7.4	0.38
Pillar	21.5	21.5	0.50
FCE	13.9	26.2	0.11
IRF	16.9	16.9	0.50
Atrophy	9.2	15.4	0.55
CNV	15.4	18.5	0.38

Abbreviations: BCVA: best corrected visual acuity; Pre-VL: pre-vitelliform; VL: vitelliform; SRF: subretinal fluid; FCE: focal choroidal excavation; IRF: intraretinal fluid; VA: visual acuity; CST: central subfield thickness; CNV: choroidal neovascularization. P-value <0.05 considered statistically significant.