

Supplementary Table 1. Background and demographics of patients who had or had not experienced adverse reactions to topical PG**Eyelash hypertrichosis**

Experienced an adverse reaction	(+) N=76		(-) N=32		p-value [†]
	Frequency	%	Frequency	%	
Age, mean±SD (years)	60.7±11.0		64.4±12.3		0.136
Sex:					0.137
Male	43	56.6	23	71.9	
Female	33	43.4	9	28.1	

Eyelid hyperpigmentation

Experienced an adverse reaction	(+) N=77		(-) N=31		p-value [†]
	Frequency	%	Frequency	%	
Age, mean±SD (years)	60.9±10.7		62.2±12.7		0.586
Sex:					0.919
Male	48	62.3	19	61.3	
Female	29	37.7	12	38.7	

DUES

Experienced an adverse reaction	(+) N=59		(-) N=47		p-value [†]
	Frequency	%	Frequency	%	
Age, mean±SD (years)	60.5±11.1		63.1±11.5		0.248
Sex:					0.182
Male	34	57.6	33	70.2	
Female	25	42.4	14	29.8	

†A two-sample *t*-test was performed to compare age of subjects with or without experiencing adverse reactions to topical PG. A chi-squared test was performed to compare the number of patients who experienced an adverse reaction by gender groups. The statistical significance of the test was set at less than 5%. PG, prostaglandin; SD, standard deviation; DUES, deepening of the upper eyelid sulcus; N, number of patients.